arXiv:1409.0371v1 [nlin.SI] 1 Sep 2014

Soliton solutions, Bäcklund transformation and Lax pair for coupled Burgers system via Bell polynomials

Ömer Ünsal*, Filiz Taşcan

Eskişehir Osmangazi University, Art-Science Faculty, Department of Mathematics-Computer, Eskişehir-TÜRKİYE

Email : ounsal@ogu.edu.tr; ftascan@ogu.edu.tr

September 2, 2014

Abstract

In this work, we apply the binary Bell polynomial approach to coupled Burgers system. In other words, we investigate possible integrability of referred system. Bilinear form and soliton solutions are obtained, some figures related to these solutions are given. We also get Bäcklund transformations in both binary Bell polynomial form and bilinear form. Based on the Bäcklund transformation, Lax pair is obtained. Namely, this is a study in which integrabilitiy of coupled burgers system is investigated.

Keywords: Bell polynomials, soliton solutions, Backlund transformation, Lax pair.

PACS (2010) : 02.30 Jr, 02.30 lk, 05.45 Yv, 11.30 -j.

1 Introduction

Nonlinear evolution equations (NLEEs) and systems of NLEEs are known to describe a wide variety of phenomena in physics, engineering, applied mathematics, chemistry, biology. In this manner, integrability of NLEEs and systems of NLEEs has been more important and topic of many papers. During the past decades, many scientific workers have presented papers which includes study of complete integrability of NLEEs. For instance, the Darboux transformation method [1], Painleve analysis method [2], inverse scattering method [3], Hirota bilinear method [4], Bäcklund transformation method [5]. Hirota bilinear method is based on the construction of a bilinear form of given NLEE. Lambert and his co-workers proposed an useful approach to study the bilinear form, bilinear Bäcklund transformation, Lax pair, Darboux transformation by using Bell polynomials [6, 7]. The Bell polynomial approach is very practical to characterize the bilinearizable equations. Fan [8] generalized this method to search the complete integrability of the nonisospectral and the variable-coefficient equations and proposed a procedure to obtain the infinite conservation laws of the NLEEs. Lambert and Springael constructed Bell polynomial approach for systems of NLEEs and NLEEs which can be transformed to a system of NLEEs. They also gave some applications [9]. Then, some authors presented more applications [10, 11, 12, 13, 14].

This paper is organized as follows: In Section 2, we briefly present some notations related to Bell polynomials used in the literature. In Section 3, bilinear formalism and soliton solutions are obtained, some figures correspond to the soliton solutions are given.

*Corresponding Author. E-mail address: ounsal@ogu.edu.tr

In section 4, Bäcklund transformation and Lax pair are costructed. In section 4, some conclusions are given.

2 Bell polynomials and binary ones

In this section, we give the basic definitions and fundamentals of multi-dimensional Bell polynomials and binary Bell polynomials. For details, please refer to [13, 14, 15, 16, 17, 18, 19, 20, 21]. Consider the function $f = f(x_1, x_2, ..., x_l)$ with multi-variables in C^∞. The multi-dimensional Bell polynomials are given as

$$\begin{array}{rcl} Y_{n_1x_1,...,n_lx_l}(f) & = & Y_{n_1,...,n_l}(f_{r_1x_1,...,r_lx_l}) \\ & = & e^{-f}\partial_{x_1}^{n_1}...\partial_{x_l}^{n_l}e^f, \end{array} \tag{1}$$

where

$$f_{r_1x_1,...,r_lx_l} = \partial_{x_1}^{r_1}...\partial_{x_l}^{r_l}f,$$

$$r_1 = 0, ..., n_1; ...; r_l = 0, ..., n_l.$$

For the simplest case $f = f(x)$, the one-dimensional Bell polynomials are expressed as

$$\begin{array}{l} Y_1 = f_x, Y_2 = f_{2x} + f_x^2, \\ Y_3 = f_{3x} + 3f_xf_{2x} + f_x^3, ... \end{array} \tag{2}$$

If function has two variables, namely for the $f = f(x, t)$, the associated two-dimensional Bell polynomials are

$$\begin{array}{rcl} Y_{x,t} & = & f_{x,t} + f_xf_t, \\ Y_{2x,t} & = & f_{2x,t} + f_{2x}f_t + 2f_{x,t}f_x + f_x^2f_t, ... \end{array} \tag{3}$$

The multi-dimensional binary Bell polynomials take the following forms

$$\mathcal{Y}_{n_1x_1,...,n_lx_l}(v, w) = Y_{n_1,...,n_l}(f) \tag{4}$$

for

$$f_{r_1x_1,...,r_lx_l} = \begin{cases} v_{r_1x_1,...,r_lx_l}, \text{ when } r_1 + ... + r_l \text{ is odd} \\ w_{r_1x_1,...,r_lx_l}, \text{ when } r_1 + ... + r_l \text{ is even.} \end{cases}$$

The first few lowest order binary Bell polynomials read

$$\begin{array}{l} \mathcal{Y}_x(v) = v_x, \ \mathcal{Y}_{2x}(v, w) = w_{2x} + v_x^2, \\ \mathcal{Y}_{x,t}(v, w) = w_{x,t} + v_xv_t, \\ \mathcal{Y}_{3x}(v, w) = v_{3x} + 3v_xw_{2x} + v_x^3, \\ \mathcal{Y}_{4x}(v, w) = w_{4x} + 3w_{2x}^2 + 4v_xv_{3x} + 6v_x^2w_{2x} + v_x^4, ... \end{array} \tag{5}$$

The link between the $\mathcal{Y}$-pollynomials and the standard Hirota bilinear formula $D_{x_1}^{n_1}...D_{x_l}^{n_l}F.G$ can be expressed by the following identity:

$$\mathcal{Y}_{n_1x_1,...,n_lx_l}(v = \ln F/G, w = \ln FG) = (FG)^{-1}\, D_{x_1}^{n_1}...D_{x_l}^{n_l}F.G, \tag{6}$$

where $n_1 + n_2 + ... + n_l \geq 1$, and operators $D_{x_1}, ..., D_{x_l}$ are the classical Hirota's bilinear operators defined by

$$D_{x_1}^{n_1}...D_{x_l}^{n_l}F.G = \left(\partial_{x_1} - \partial_{x_1'}\right)^{n_1} ... \left(\partial_{x_l} - \partial_{x_l'}\right)^{n_l} F(x_1, ..., x_l) \times G(x_1', ..., x_l') \mid_{x_1'=x_1,...,x_l'=x_l}.$$

Furthermore, the relationship between the binary Bell polynomials and Lax Pair by use of the Hopf-Cole transformation is given by the expression

$$\mathcal{Y}_{n_1x_1,n_2x_2}(v = \ln\psi, w = v + Q) = \psi^{-1}\sum_{r=0}^{n_1}\sum_{s=0}^{n_2}\binom{n_1}{r}\binom{n_2}{s}\mathcal{Y}_{rx_1,sx_2}(0, Q)\,\partial_{x_1}^{n_1-r}\partial_{x_2}^{n_2-s}\psi, \tag{7}$$

where ψ and Q are both the functions of x_1 and x_2.

3 Binary Bell polynomial form and soliton solutions

Now we consider the coupled Burgers system [22]

$$u_t - 2uu_x - v_{xx} = 0 \tag{8a}$$

$$v_{yt} - u_{xxy} - 2uv_{xy} - 2u_x v_y = 0. \tag{8b}$$

Wang et al [22] obtained infinitely many generalized symmetries of system (8). We introduce two dimensionless fields p and q which are the functions of x, y and t. Taking the transformation

$$u = p_x \ , \ v = q_x \tag{9}$$

into account and integrating the resulting equation once with respect to x, we get the Eq.(8a) as

$$p_t - p_x^2 - q_{xx} = 0. \tag{10}$$

Substituting (9) into Eq.(8b), using Eq.(10) and integrating the resulting equation once with respect to x, we obtain

$$q_{yt} + p_y p_t - p_y \left(p_x^2 + q_{xx}\right) - p_{xxy} - 2p_x q_{xy} = 0. \tag{11}$$

From Eqs.(10) and (11) we find binary Bell polynomial form of the system (8) as

$$\mathcal{Y}_t(p,q) - \mathcal{Y}_{xx}(p,q) = 0 \tag{12a}$$

$$\mathcal{Y}_{yt}(p,q) - \mathcal{Y}_{xxy}(p,q) = 0. \tag{12b}$$

Through expression (6) and setting that $p = \ln(f/g)$, $q = \ln(fg)$ we get bilinear form of system (8) as

$$\left[D_t - D_x^2\right] f.g = 0 \tag{13a}$$

$$\left[D_y D_t - D_x^2 D_y\right] f.g = 0 \tag{13b}$$

where f and g are the functions of x, y and t.

As follows, the Eqs.(13) can be easily solved to obtain the multi-soliton solutions of system (8) by using the Hirota's bilinear method.

Let f and g be functions in the form

$$f = \epsilon f_1 + \epsilon^2 f_2 + ... \tag{14a}$$

$$g = 1 + \epsilon g_1 + \epsilon^2 g_2 + ... \tag{14b}$$

Substituting expressions (14) into Eqs. (13) and collecting the coefficients of the same power of ϵ, we have

$$\epsilon^1 : \ \left[D_t - D_x^2\right](f_1.1) = 0 \tag{15a}$$

$$\left[D_y D_t - D_x^2 D_y\right](f_1.1) = 0 \tag{15b}$$

$$\epsilon^2 : \ \left[D_t - D_x^2\right](f_1.g_1 + f_2.1) = 0 \tag{16a}$$

$$\left[D_y D_t - D_x^2 D_y\right](f_1.g_1 + f_2.1) = 0 \tag{16b}$$

$$\epsilon^3 : \ \left[D_t - D_x^2\right](f_1.g_2 + f_2.g_1 + f_3.1) = 0 \tag{17a}$$

$$\left[D_y D_t - D_x^2 D_y\right](f_1.g_2 + f_2.g_1 + f_3.1) = 0 \tag{17b}$$

$$\vdots$$

(i) One-Soliton Solution

In order to find one soliton solution of system (8), we take

$$f_1 = ae^{\xi},\ g_1 = be^{\xi},\ \xi = kx + my + wt,$$
$$f_i(x,y,t) = 0, g_i(x,y,t) = 0\ i = 2,3,4,... \tag{18}$$

where a, b, k and m are all arbitrary nonzero real constants. w is a real constant to be determined. Solving (15)-(17) which found by substituting expressions (14) into (13), we get $w = k^2$. So the one soliton solutions for system (8) is obtained in the form

$$u = \frac{k}{1 + be^{kx+my+k^2t}} \tag{19a}$$

$$v = \frac{k\left(1 + 2be^{kx+my+k^2t}\right)}{1 + be^{kx+my+k^2t}} \tag{19b}$$

(i) Two-Soliton Solution

Similarly, we look for the two-soliton solution by choosing

$$f_1 = a_1e^{\xi_1} + a_2e^{\xi_2},\ g_1 = b_1e^{\xi_1} + b_2e^{\xi_2},$$
$$f_2 = a_3e^{\xi_1+\xi_2},\ g_2 = b_3e^{\xi_1+\xi_2},\ \xi_i = k_ix + m_iy + w_it,\ i = 1,2 \tag{20}$$

where $a_1, a_2, a_3, b_1, b_2, b_3, k_1, k_2$ are all arbitrary nonzero constants. We truncate the expressions (14) to $f_2(x,y,t)$ and $g_2(x,y,t)$, respectively. Then, corresponding two-soliton solution is

$$u = \left[\ln\left(\frac{a_1e^{\xi_1} + a_2e^{\xi_2} + a_3e^{\xi_1+\xi_2}}{1 + b_1e^{\xi_1} + b_2e^{\xi_2} + b_3e^{\xi_1+\xi_2}}\right)\right]_x \tag{21a}$$

$$v = \left[\ln\left(\left(a_1e^{\xi_1} + a_2e^{\xi_2} + a_3e^{\xi_1+\xi_2}\right)\left(1 + b_1e^{\xi_1} + b_2e^{\xi_2} + b_3e^{\xi_1+\xi_2}\right)\right)\right]_x \tag{21b}$$

where

$$a_1 = \frac{m_1k_1a_3}{b_2(k_1-k_2)(m_1-m_2)}\ ,\ b_1 = \frac{m_2k_2a_3}{a_2(k_1-k_2)(m_1-m_2)}\ ,\ w_i = k_i^2. \tag{22}$$

Some figures which corresponds to soliton solutions found above are given below. Figs. 1-2 and 3-6 represents (19) and (21) respectively. Figs. 1 and 2 show the motion of the shock waves. Fig. 3 shows head on collision of two kink type solitary waves. The collision is inelastic, that's why the shape of waves is changed after the collision. Fig. 5 depicts the interaction of two bell shaped solitary waves. Waves move emerging into one after the collision. Then the collision is inelastic. Figs. 4 and 6 show the interaction of two kink type solitons. The solitons keep their own sahpe and direction after the interaction. So collision is elastic.



u:Fig. 1. $t = 0, k = -0.6, m = 0.2, b = 0.4$



v:Fig. 2. $t = 0, k = -0.1, m = 1.5, b = 0.1$



u:Fig. 3. $t = 0, k_1 = -3, k_2$=2, $m_1 = 0.5, m_2 = -2, a_2 = -0.2, a_3 = -0.2,$ $b_2 = 0.3, b_3 = 1$



v:Fig. 4. $t = 0, k_1 = -3, k_2 = 2,$ $m_1 = 0.5, m_2 = -2, a_2 = -0.2, a_3 = -0.2,$ $b_2 = 0.3, b_3 = 1$



u:Fig. 5. $t = 0, k_1 = -0.1, k_2$ =0.8, $m_1 = 1.2, m_2 = -0.3, a_2 = -0.2, a_3 = -0.2,$ $b_2 = 0.1, b_3 = 1$



v:Fig. 6. $t = 0, k_1 = -0.1, k_2 = 0.8,$ $m_1 = 1.2, m_2 = -0.3, a_2 = -0.2, a_3 = -0.2,$ $b_2 = 0.1, b_3 = 1$

4 Bäcklund transformation and Lax Pair

In this section, we get the bilinear Bäcklund transformation in both the binary Bell polynomial and bilinear forms. We also obtain Lax pair of system (8). We take

$$P_1 = [\mathcal{Y}_t(p', q') - \mathcal{Y}_{xx}(p', q')] - [\mathcal{Y}_t(p, q) - \mathcal{Y}_{xx}(p, q)] \tag{23}$$

$$P_2 = [\mathcal{Y}_{yt}(p', q') - \mathcal{Y}_{xxy}(p', q')] - [\mathcal{Y}_{yt}(p, q) - \mathcal{Y}_{xxy}(p, q)] \tag{24}$$

where (p', q') and (p, q) satisfy the Eqs. (12). Supposing that $p' = \ln(f'/g')$ and $q' = \ln(f'g')$, using the relations

$$\begin{array}{llll} v_1 = \ln(g'/g), & v_2 = \ln(f'/f), & v_3 = \ln(f'/g), & v_4 = \ln(g'/f), \\ w_1 = \ln(g'g), & w_2 = \ln(f'f), & w_3 = \ln(f'g), & w_4 = \ln(g'f), \end{array} \tag{25}$$

$$\begin{array}{ll} p' - p = v_2 - v_1 = w_3 - w_4 & p' + p = v_3 - v_4 = w_2 - w_1 \\ q' - q = v_1 + v_2 = v_3 + v_4 & q' + q = w_1 + w_2 = w_3 + w_4 \end{array} \tag{26}$$

we obtain the binary Bell polynomial form of Bäcklund transformation for the system (8)

$$\begin{array}{l} \mathcal{Y}_t(v_1, w_1) + \mathcal{Y}_{xx}(v_1, w_1) = 0 \\ \mathcal{Y}_t(v_2, w_2) + \mathcal{Y}_{xx}(v_2, w_2) = 0 \\ \mathcal{Y}_x(v_3, w_3) - \mu e^{v_1 - v_2} = 0 \\ \mathcal{Y}_{xy}(v_3, w_3) - d\mathcal{Y}_x(v_3, w_3) + \mu e^{v_1 - v_2} \mathcal{Y}_y(v_4, w_4) = 0 \end{array} \tag{27}$$

where μ and d are arbitrary real constants. With the aid of (6), we can also express Bäcklund transformation in the bilinear form as follows:

$$\begin{array}{l} \left[D_t + D_x^2\right] g'.g = 0 \\ \left[D_t + D_x^2\right] f'.f = 0 \\ D_x f'.g - \mu f' g = 0 \\ \left[D_x D_y - d D_x\right] f'.g + \mu D_y g'.f = 0. \end{array} \tag{28}$$

As a result of the relations (25)-(26), relations

$$v_2 = v_3 - p, \quad w_2 = w_3 + p, \quad v_4 = v_1 - p, \quad w_4 = w_1 + p \tag{29}$$

can be written [9]. Taking that

$$w_i = v_i + Q_i, \quad v_i = \ln \psi_i, \quad (i = 1, 3) \tag{30}$$

we can use the relations

$$Q = w_1 - v_1 = w_3 - v_3 = Q_1 = Q_3. \tag{31}$$

Using (29)-(31), we find the Lax pair for the system (8) in the form

$$\begin{array}{c} \psi_{1,y} = \frac{d}{2}\psi_1 - \frac{e^{-p}(q_{xy} - p_{xy})}{2\mu}\psi_3 \\ \psi_{1,t} = \left(p_{xx} - q_{xx}\right)\psi_1 - \psi_{1,xx} \\ \psi_{3,x} = \mu e^p \psi_1 \\ \psi_{3,t} = \mu p_x e^p \psi_1 - \mu e^p \psi_{1,x}. \end{array} \tag{32}$$

Compatibility conditions $\psi_{1,yt} = \psi_{1,ty}$, $\psi_{3,xt} = \psi_{3,tx}$ can easily be verified.

5 Conclusion

In this study, we have investigated the coupled Burgers system with the aid of Bell polynomials. In this sense, we have obtained Bilinear form. Using bilinear form we have gotten soliton solutions and discussed these solutions analytically. Some figures of one-soliton and two-soliton solutions have been given. Bäcklund transformations in both binary Bell polynomial form and bilinear form have been given. Then, Lax pair has been constructed. We hope that the present findings may be useful in further works.

References

[1] Matveev, V.B., Salle, M.A., Darboux transformation and solutions, Springer, Berlin (1980).

[2] Li, X.N., Wei, G.M., Liang, Y.Q., Painleve analysis and new analytic solutions for variable-coefficient Kadomtsev-Petviashvili equation with symbolic computation, Appl. MAth. Comput. 216, (2010) 3568.

[3] Ablowitz, M.J., Clarkson, P.A., Solitonsi nonlinear evolution equations and inverse scattering, Cambridge University Press, Cambridge (1991).

[4] Wazwaz, A.M., Multiple-soliton solutions for the KP equation by Hirota's bilinear method and by the tanh–coth method, Applied Mathematics and Computation, 190,1 (2007) 633.

[5] Miura, M.R., Backlund Transformation, Springer-Verlag, Berlin, (1978).

[6] Gilson, C., Lambert, F., Nimmo, J., Willox, R., On the combinatorics of the Hirota D-operators, Proc. R. Soc. Lond. A, 452, (1996) 223.

[7] Lambert, F., Springael, J., Soliton equations and simple combinatorics, Acta. Appl. Math., 102, (2008) 147.

[8] Fan, E., The integrability of nonisospectral and variable-coefficient KdV equation with binary Bell polynomials, Physics Letters A, 375, (2011) 493.

[9] Lambert, F., Springael, J., On a direct procedure for the disclosure of Lax pairs and Bäcklund transformations, Chaos, Solitons and Fractals, 12, (2001) 2821.

[10] Sun, Y., Tian, B., Sun, W.R., Jiang, Yan., Wang, Y.P., Huang, Z.R., Bäcklund transformation and N-soliton solutions for a (2+1)-dimensional nonlinear evolution equation in nonlinear water waves.

[11] Qin, Y., Gao, Y.T., Yu, X., Meng, G.Q., Bell polynomial approach and N-Soliton Solutions for a Coupled KdV-mKdV System, Commun. Theor. Phys., 58, (2012) 73.

[12] Wang, Y.F., Tian, B., Wang, P., Li, M., Jiang, Y., Bell-polynomial approach and soliton solutions for the Zhiber-Shabat equation and (2+1)-dimensional Gardner equation with symbolic computation, Nonlinear Dyn., 69, (2012) 2031.

[13] Wang, P., Tian, B., Liu, W.J., Lü, X., Jiang, Y., Lax pair, Bäcklund transformation and multi-soliton solutions for the Boussinesq-Burgers equations from shallow water waves, Applied Mathematics and Computation, 218, (2011) 1726.

[14] Jiang, Y., Tian, B., Liu, W.J., Sun, K., Li, M., Soliton solutions and integrability for the generalized variable-coefficient extended Korteweg-de Vries equation in fluids, Applied Mathematics Letters, 26, (2013) 402.

[15] Bell, E.T., Exponential polynomials, Ann. Math. 35, (1934) 258.

[16] Lambert, F., Loris, I., Springael, J., Willox, R., On modified NLS, Kaup and NLBqequations: differential transformations and bilinearization, J. Phys. A: Math. Gen. 27, (1994) 8705..

[17] Lambert, F., Springael, J., Construction of Bäcklund transformations with binary Bell polynomials, J. Phys. Soc. Jpn., 66, (1997) 2211.

[18] Lambert, F., Loris, I., Springael, J., Classical Darboux transformations and the KP hierarchy, Inverse Problems, 17, (2001) 1067.

[19] Lambert, F., Leble, S., Springael, J., Binary Bell polynomials and Darboux covariant Lax pairs, Glasg. Math. J.,43A, (2001) 53.

[20] Luo, L., New exact solutions and Backlund transformation for Boiti-Leon-Manna-Pempinelli equation, Physics Letters A, 375, (2011) 1059.

[21] Jiang, Y., Tian, B., Liu, W.J., Li, M., Wang, P., Sun, K., Solitons, Backlund transformation, and Lax pair for the (2+1)-dimensional Boiti-Leon-Pempinelli equation for the water waves, Journal of Mathematical Physics, 51, (2010) 093519.

[22] Wang, Y.J., Liang, Z.F., Tang, X.Y., Infinitely many generalized symmetries and Painleve analysis of a (2+1)-dimensional Burgers system, Physica Scripta, 89, (2014) 025201.